The AAO FORUM FOR OSTEOPATHIC THOUGHT JOURNAL

Official Publication of the American Academy of Osteopathy.

TRADITION SHAPES THE FUTURE

Volume 26 • Number 3 • September 2016



The pilot study that begins on page 21 was designed to investigate whether using a shim to level the iliac crests changes the diagnostic reliability of the standing flexion test and to appraise interexaminer reliability.

The American Academy of Osteopathy is your voice...

in teaching, promoting, and researching the science, art, and philosophy of osteopathic medicine, with the goal of integrating osteopathic principles and osteopathic manipulative treatment in patient care.

If you are not already a member of the American Academy of Osteopathy (AAO), the AAO Membership Committee invites you to join the Academy as a 2016-17 member. The AAO is your professional organization. It fosters the core principles that led you to become a doctor of osteopathic medicine.

For \$5.54 a week (less than the price of a large specialty coffee at your favorite coffee shop) or just 79 cents a day (less than the cost of a bottle of water), you can become a member of the professional specialty organization dedicated to you and osteopathic manipulative medicine (OMM).

Your membership dues provide you with:

- a national advocate for OMM, both within the profession and with health care policy-makers and third-party payers.
- a champion that is monitoring closely and responding rapidly to the standards being developed for the single accreditation system for graduate medical education.
- referrals of patients through the "Find a Physician" tool both on the AAO website and at the FindOMM.org URL, as well as from calls to the AAO office.
- discounts on continuing medical education at the AAO's annual Convocation and its weekend courses.
- automatic acceptance of AAO-sponsored courses by the American Osteopathic Board of Neuromusculoskeletal Medicine, the only certifying board for manual medicine in the world today.

- networking opportunities with peers.
- discounts on books in the AAO's online store.
- complimentary subscription to *The AAO Journal*, published electronically 4 times annually.
- complimentary subscription to the online *AAO Member News*, published 8 times annually.
- weekly *OsteoBlast* e-newsletters, featuring research on manual medicine from peer-reviewed journals around the world.
- practice promotion materials, such as the AAO-supported "American Health Front!" segment on OMM.
- discounts on advertising in AAO publications and in materials for the AAO's Convocation.
- the fellow designation of FAAO, which recognizes DOs for promoting OMM through teaching, writing, and professional service and which is the only earned fellowship in the osteopathic medical profession.
- promotion and grant support of research on the efficacy of OMM.
- support for the future of the profession through the Student American Academy of Osteopathy, the National Undergraduate Fellows Association, and the Postgraduate American Academy of Osteopathy.

If you have any questions regarding membership or membership renewal, contact AAO Membership Liaison Susan Lightle at SLightle@academyofosteopathy.org or at (317) 879-1881, ext. 217.



Georgia Campus-Philadelphia College of Osteopathic Medicine is located in Suwanee, GA in Gwinnett County. We're located in the bustling metro Atlanta region, close to the North Georgia mountains, lakes, downtown cultural amenities, and professional sporting events. We have the best of big-city vibrancy and small-town community. We are currently seeking qualified individuals for a full time faculty position in the Department of Osteopathic Manipulative Medicine (OMM).

Faculty - Osteopathic Manipulative Medicine

Individual will be expected to teach osteopathic medical students in both lecture and laboratory sessions in all four years of the curriculum,

see patients and develop an outpatient clinic for M-3 month long rotation, plan and supervise OMM Inpatient Student Service, assist in preparation of OMM video clips and tutorials, participate in existing research and initiate new OMM research, assist in planning and production of new publications, and assist in planning and supervision of the OMM Residency.

The successful applicant will have a D.O. degree and proficiency in osteopathic manipulative medicine. The candidate needs to have or be eligible for a license to practice Osteopathic Medicine in the State of Georgia. Board Certification or eligibility by the AOBNMM or AOBSPOMM is required. Additional Board Certification or eligibility by the AOBFM is desirable

To apply for the position, send via E-mail a personal statement describing interest in and qualifications for this position, a curriculum vitae, and names and addresses of three references, preferably from current or former supervisors.

Department of Human Resources, 625 Old Peachtree Road, Suwanee, GA 30024 Phone **(678)225-7500**, Fax **(678)225-7519**, E-mail: **hr@pcom.edu**





3500 DePauw Blvd, Suite 1100 Indianapolis, IN 46268-1136 (317) 879-1881 • fax: (317) 879-0563 editoraaoj@gmail.com www.academyofosteopathy.org

The AAO Journal

Brian E. Kaufman, DO, FACOI, FACP Scientific editor Katherine A. Worden, DO, MS Associate editor Raymond J. Hruby, DO, FAAODist ... Scientific editor emeritus Lauren Good Managing editor

AAO Publications Committee

Hollis H. King, DO, PhD,	Brian E. Kaufman, DO,
FAAO, chair	FACOI, FACP
William J. Garrity, DO,	Bobby Nourani, DO
vice chair	Hallie J. Robbins, DO
Claire M. Galin, DO	Katherine A. Worden, DO, MS
Edward Keim Goering, DO	Paul R. Rennie, DO, FAAO,
Stephen I. Goldman, DO,	Board of Trustees' liaison
FAAO, FAOASM	Michael Smith, OMS III,
Raymond J. Hruby, DO, MS,	SAAO liaison
FAAODist	

American Academy of Osteopathy

Laura E. Griffin, DO, FAAO	President
Michael P. Rowane, DO, MS, FAAO, FAAFP Pres	ident-elect
Sherri L. Quarles	ve director

The AAO Journal is the official publication of the American Academy of Osteopathy. Issues are published 4 times a year.

The AAO Journal is not responsible for statements made by any contributor. Opinions expressed in *The AAO Journal* are those of the authors and do not necessarily reflect viewpoints of the editors or official policy of the American Academy of Osteopathy or the institutions with which the authors are affiliated, unless specified.

Although all advertising is expected to conform to ethical medical standards, acceptance does not imply endorsement by this journal or by the American Academy of Osteopathy.

Subscription rate for AAO nonmembers: \$60 per year. To subscribe, contact AAO Communications Specialist Lauren Good at LGood@academyofosteopathy.org.

Send all address changes to SLightle@academyofosteopathy.org.

ISSN 2375-5717 (online) ISSN 2375-5776 (print)



The mission of the American Academy of Osteopathy is to teach, advocate, and research the science, art, and philosophy of osteopathic medicine, emphasizing the integration of osteopathic principles, practices, and manipulative treatment in patient care.

Editorial

View From the Pyramids:
Reframing the "Marijuana Menace"5
Brian E. Kaufman, DO, FACOI, FACP

ORIGINAL CONTRIBUTIONS

The Incidence of Somatic Dysfunction in Patients
With Sudden Onset Atraumatic Neck Pain:
A Retrospective Case Note Study9
Karen Teten Snider, DO, FAAO; Amrien Ghouse, OMS VI; Sheri
Shiyi He, BS; and Vanessa K. Pazdernik, MS
Does Using a Shim Improve the Diagnostic Reliability of the
Standing Flexion Test? A Pilot Study
Alexa Paulina Marquez, OMS III, MS; Michelle Yim, OMS
IV, MPH; Biana Zelimkhanian, OMS IV, MS; John Quiamas,
OMS IV, MS; Rachel E. Davidge, DO; Matthew Siri, DO; and
Raymond J. Hruby, DO, MS, FAAODist

REGULAR FEATURES

AAO Calendar of Events	4
CME Certification of Home Study	20
Component Society Calendar of Events	

The advertising rates listed below are for *The AAO Journal*, the official peerreviewed publication of the American Academy of Osteopathy (AAO). AAO members and AAO component societies are entitled to a 20% discount on advertising in this journal. Call the AAO at (317) 879-1881, ext. 211, for more information.

2016 Advertising Rates per Placement				
	Placed 1 time	Placed 2 times	Placed 4 times	
Full page (7.5" x 9.5")	\$600	\$570	\$540	
One-half page (7.5" x 4.5")	\$400	\$380	\$360	
One-third page (2.25" x 9.5")	\$300	\$285	\$270	
Quarter page (3.5" x 4.5")	\$200	\$190	\$180	
Classified	\$1 per 7 characters, spaces not included			



AAO Calendar of Events



Mark your calendar for these upcoming Academy meetings and educational courses.

2016-2017

Sept. 15	AAO Board of Trustees' meeting—9 a.m. to 5 p.m. Pacific—Anaheim, California	Jan. 2	AAO office closed in observance of New Year's Day
Sept. 16	AAO Leadership Forum—9 a.m. to 3 p.m. Pacific—Anaheim, California	Jan. 11	Committee on Fellowship in the AAO's teleconference—8:30 p.m. Eastern
Sept. 17	AAO Education Committee's meeting—3 to 4:30 p.m. Pacific—Anaheim, California	Jan. 20-22	Osteopathic Management of Chronic Pain: Addressing Chronic Fatigue Syndrome, Eibrogradia Multinla Salaragia and
Sept. 17-20	AAO at OMED: Osteopathic Neuromusculoskeletal Medicine in the 21st Century—Daniel G. Williams, DO, program chair—Anaheim (California) Convention Center		Neuroinflammation—Bruno J. Chikly, MD, DO (France), course director—Midwestern University/Arizona College of Osteopathic Medicine in Glendale
Oct. 15	Committee on Fellowship in the AAO's meeting—AAO office in Indianapolis	Feb. 3-4	AAO Education Committee's meeting—AAO office in Indianapolis
Oct. 21-23	What's the Point? Multifaceted Clinical Approaches to Viscerosomatic Reflexes— Michael L. Kuchera, DO, FAAO, course director: William H. Devine, DO, and Richard	March 18-21	Pre-Convocation—The Strategic Crossroads of the Body—Jean-Pierre Barral, DO (France), featured speaker—The Broadmoor, Colorado Springs, Colorado
	A. Feely, DO, FAAO, FCA, course faculty —Midwestern University/Arizona College of Osteopathic Medicine in Glendale	March 19-21	Pre-Convocation—Brain 1: Palpating and Treating the Brain, Brain Nuclei, White Matter and Spinal Cord—Bruno J. Chikly, MD, DO (France), course director—The Broadmoor,
Nov. 24	Thanksgiving Day—AAO office closed		Colorado Springs, Colorado
Dec. 1	FAAO applications due	March 19-21	Pre-Convocation—Fascial Distortion Model: Avial Spine Todd A Capitrant DO MHA
Dec. 2-4	Fulford's Advanced Percussion Hammer— Richard W. Koss, DO, course director— University of North Texas Health Science Center		course director—The Broadmoor, Colorado Springs, Colorado
	Texas College of Osteopathic Medicine in Fort Worth	March 22	Pre-Convocation—Cervical Glymphatics— Frank Willard, PhD, course faculty—The Breadmoor Colorado Springs Colorado
Dec. 2-4	Cranial Approach of Beryl E. Arbuckle, DO-		broadmoor, Colorado Springs, Colorado
	Kenneth J. Lossing, DO, course director— Midwestern University/Arizona College of Osteopathic Medicine in Glendale	March 22-27	AAO Convocation—The Balance Point: Bringing the Science and Art of Osteopathic Medicine Together—Natalie Ann Nevins, DO,
Dec. 20	Committee on Fellowship in the AAO's teleconference—8:30 p.m. Eastern		program chair—1ne broadmoor, Colorado Springs, Colorado
Dec. 26	AAO office closed in observance of Christmas Day	March 27	Post-Convocation—Program Directors Workshop—Eric Hunter Sharp, DO, course director—The Broadmoor, Colorado Springs, Colorado



View From the Pyramids: Reframing the "Marijuana Menace"

AAOJ Scientific Editor Brian E. Kaufman, DO, FACOI, FACP

EDITORIAL

The use of cannabis to treat medical conditions is established in documents dating back thousands of years. Cannabis has been used in the United States since the 1600s as hemp production was encouraged and needed for industry.¹ The use of the leaf for recreation is generally attributed to the influx of Mexican immigrants in the early third of the twentieth century. Fear of the immigrants, along with frustrations from the effects of the Great Depression resulted in backlash against the "Marijuana Menace."¹ This instigated a flurry of research which linked the use of cannabis with violent crimes and socially deviant behaviors, committed by the so-dubbed "racially inferior" communities.¹

These fear-based, and largely content-devoid, assertions led to a number of progressively more restrictive laws limiting and then eliminating the use of cannabis. From the passage of the Marijuana Tax Act of 1937 until now, the use of cannabis has been a criminal offense under both federal and state laws.¹ Throughout the years, the degree of severity has waxed and waned. On November 30, 2011, the governors of 2 states petitioned the US Drug Enforcement Administration to change the status of medical cannabis,² and that brings us to the present.

Since learning the above, I will not use the term *medical marijuana* but instead use the term *medical cannabis* so as not to perpetuate

"

The illegality of cannabis is outrageous, an impediment to full utilization of a drug which helps produce the serenity and insight, sensitivity and fellowship so desperately needed in this increasingly mad and dangerous world.

-Carl Sagan⁹

Opinions expressed in *The AAO Journal* are those of the authors and do not necessarily reflect viewpoints or official policy of the American Academy of Osteopathy.

the fear and biased terminology that harkens to the "Mexican Menace."

On August 11, 2016, the DEA announced that they denied the petition to reclassify cannabis under the controlled substance act and thereby determined that cannabis will remain a Schedule 1 drug. By definition,³ Schedule 1 drugs

- have a high potential for abuse
- have no currently accepted medical uses in the US
- lack accepted safety for using the drug under medical supervision

The decision to not reclassify cannabis was a disappointment to both patients and physicians. Cannabis has been shown to be helpful for many conditions, from epilepsy to chronic pain. Medical cannabis has also been shown to decrease the need for opiates, which in the present climate is extremely useful.⁴

As with all substances physicians prescribe, there are contraindications and cautions, and appropriate risk-benefit analysis should always take place. However, cannabis has an extremely good safety profile when compared to many of the medications we routinely prescribe. Nonsteroidal anti-inflammatories carry a risk of death with acute use, and in the US, NSAIDs account for at least 16,500 deaths a year.⁵ Add to that the deaths attributed to Coumadin and the other novel anticoagulants, to antipsychotic medications, and to many others we prescribe daily, and cannabis does not come close.

A 2014 study reported 2 cases of death from acute use of cannabinoids, attributed to cardiovascular complications.⁶ Besides that, the risks are all potentially with chronic use, and they are related to smoking, not the actual substance. There can also be secondary contributions, such as in use in a patient with schizophrenia trig-

(continued on page 6)

(continued from page 5)

gering psychosis. The California Academy of Addiction Medicine (CSAM), states the following:

The fact that marijuana is clearly addictive for a minority of people, and especially so for adolescents and children, does not mean that the public does not have a right to legalize its use. We currently permit the marketing, sale and consumption of multiple legal, though addictive, substances—eg, caffeine, nicotine and alcohol. We also permit the prescription medical use of very powerfully addictive drugs—eg, tranquilizers (benzodiazepines), sleep aids, amphetamine, opiate analgesics.⁷

Given the overwhelming safety profile of cannabis, regardless of proven clinical efficacy, I find the recent determination by the DEA both shortsighted and cowardly. This is a slap in the face to all patients who depend on this as a treatment and to those physicians who recommend cannabis for their patients. Indeed, just a week after the DEA announcement, a federal court ruled that the U.S. Department of Justice cannot spend any federal monies to prosecute cannabis-related offenses that don't violate state laws.⁸ This appears to mitigate the effect of the DEA's determination, at least in part.

We have entered a time when we as physicians are having our prescribing abilities, not just over-regulated, but over-legislated, and our hands are becoming increasingly tied. In addition, we find our mouths are increasingly silenced. In the last few months, I have been told to stop prescribing opiates and to stop performing osteopathic manipulative treatment, and the federal government (via the Centers for Disease Control and Prevention) has provided a document that purports to be aimed at treating patients with chronic pain but instead seems aimed at convincing me that I am the cause of all of the problems with addiction in our society. As this is occurring, the federal government and the DEA are continuing to limit access to a potential treatment which is safer than ibuprofen.

Politics and outdated views governed by fear and ignorance have never had a place in medicine, yet they continue to be pushed into the medical space. It is incumbent on all of us to reach out to our legislators, colleagues, and national and state associations to voice our opinions.

Treating patients is our job, and we need to fight to retain our place in medical decision-making. Each day it is up to us to determine whether the safety and efficacy of a treatment is appropriate for our patients. If we wait too long, we may find that those rights and privileges that we achieved through our tenacity and sweat have been assigned elsewhere. Legalizing marijuana would make a lot of sense, I don't think there's a single case of marijuana overdose on record and tens of millions of users. It's much less dangerous than alcohol, for example.

-Noam Chomsky¹⁰

References

- Marijuana Timeline. Frontline website. http://www.pbs.org/wgbh/ pages/frontline/shows/dope/etc/cron.html. Accessed August 21, 2016.
- Memoli M. Governors petition DEA to reclassify medical marijuana. *Los Angeles Times*. November 30, 2011. http://articles.latimes. com/2011/nov/30/news/la-pn-governors-marijuana-20111130. Accessed August 29, 2016.
- 21 U.S. Code S 812 Schedules of controlled substances. Cornell University Law School Legal Information Institute website. https:// www.law.cornell.edu/uscode/text/21/812. Accessed August 29, 2016.
- Gnagey LT. Medical marijuana reduces use of opioid pain meds, decreases risk for some with chronic pain. University of Michigan website. http://ns.umich.edu/new/releases/23622-medical-marijuanareduces-use-of-opioid-pain-meds-decreases-risk-for-some-withchronic-pain. Published March 22, 2016. Accessed August 24, 2016.
- Krueger C. Ask the expert: Do NSAIDs cause more deaths than opioids? Practical Pain Management website. http://www.practicalpain-management.com/treatments/pharmacological/opioids/ask-expert-do-nsaids-cause-more-deaths-opioids. Published December 1, 2013. Updated June 23, 2016. Accessed August 29, 2016.
- Stampler L. Study claims to find first two deaths caused by marijuana. *Time* website. Published February 27, 2014. Accessed August 29, 2016.
- Marijuana's addictive potential (for healthcare professionals). California Society of Addiction Medicine website. http://www.csam-asam. org/marijuanas-addictive-potential-healthcare-professionals. Accessed August 29, 2016.
- Volokh E. Ninth Circuit bars federal prosecutions for state-lawauthorized medical marijuana. *The Washington Post* website. https:// www.washingtonpost.com/news/volokh-conspiracy/wp/2016/08/16/ ninth-circuit-bars-federal-prosecutions-for-state-law-authorized-medical-marijuana/?utm_term=.9e6caa92d7b6. Accessed August 29, 2016.
- 9. Mr. X by Carl Sagan. Dr. Lester Grinspoon's Marijuana Uses website. http://marijuana-uses.com/mr-x/. Accessed August 29, 2016.
- Cardenas LF. Noam Chomsky: Drug cartels and the growing border war. *Guernica* website. https://www.guernicamag.com/daily/luis_fernando_c225rdenas_noam/. Published August 15, 2011. Accessed August 29, 2016. ■



GET INVOLVED

The AAO Journal needs your talents.









The AAOJ needs...

- reviewers to participate in blinded peer reviews.
- osteopathic clinicians to write up osteopathic techniques, tips or variations they employ that may be published in a developing feature section of the AAOJ.
- specialists to develop osteopathic responses to articles published in nonosteopathic literature. For example, how would the osteopathic approach contribute to studies published in other scientific journals or featured in news articles?

For more information, email AAOJ Editor-in-Chief Brian E. Kaufman, DO, FACOI, FACP, at editoraaoj@gmail.com.

WHAT'S THE POINT?

MULTIFACETED CLINICAL APPROACHES TO VISCEROSOMATIC REFLEXES

Oct. 21-23, 2016 • Midwestern University/Arizona College of Osteopathic Medicine in Glendale

Palpable somatic "points" provide the osteopathic clinician with key insights directing differential diagnoses and focused approaches to patient systemic complaints. Interestingly, from those using Chapman's reflexes or Jones' counterstrain points to those who integrate acupuncture concepts or Travell's myofascial trigger points, insightful clinicians have noted a significant overlap and a great deal of similarity in clinical presentations.

Working from Chapman's approach, this course will compare and contrast a number of clinically relevant points, focus on practical diagnostic and therapeutic clues and organize the information in a practical manner for patient care. Emphasis will be placed on developing and expanding skills and strategies to speed diagnosis and recovery.

Residents, residency trainers and directors of medical education will be accorded special tips for maximizing the integrative focus of this course.

Course Faculty

Led by course director Michael L. Kuchera, DO, FAAO, the faculty will include William H. Devine, DO, and Richard A. Feely, DO, FAAO, FCA. Additional faculty may be added as attendance requires.

Course Times

Friday and Saturday, 8 a.m. to 5:30 p.m. Sunday, 8 a.m. to noon.

Continuing Medical Education

20 credits of NMM-specific AOA Category 1-A CME anticipated.

Meal Information

Breakfast will be provided each day. Lunch will be provided Friday and Saturday. Contact the AAO Event Planner Gennie Watts with special dietary needs at (317) 879-1881, ext. 220, or GWatts@ academyofosteopathy.org.

Course Location

Midwestern University/Arizona College of Osteopathic Medicine Agave Hall, OMT Lab 101, 19555 N. 59th Ave., Glendale, AZ 85308

Travel Arrangements

Contact Tina Callahan of Globally Yours Travel at (800) 274-5975 or globallyyourstravel@cox.net.

Registration Form

What's the Point? Multifaceted Clinical Approaches to Viscerosomatic Dysfunction

Oct. 21-23, 2016

Name:	AOA No.:
Nickname for badge:	
Street address:	
City:	State: ZIP:
Phone:	Fax:
Email:	

Course Faculty

Michael L. Kuchera, DO, FAAO, is a professor of osteopathic manipulative medicine at the Marian University College of Osteopathic Medicine in Indianapolis. Dr. Kuchera frequently lectures on the diagnosis and management of somatic dysfunction in patients with pain, systemic disorders or specific neuromusculoskeletal complaints.



An internationally recognized educator, **William H. Devine, DO**, is a professor of osteopathic manipulative medicine (OMM) at the Midwestern University/Arizona College of Osteopathic Medicine (MWU/AZCOM) in Glendale, where he also serves as the director of postgraduate OMM, the director of the musculoskeletal medicine residency and the coordinator of the osteopathic specialty clinic.

Since 2000, **Richard A. Feely, DO, FAAO, FCA,** has operated the Feely Center for Optimal Health in Olympia Fields, Illinois, where he uses osteopathic manipulation and acupuncture to provide relief from headaches and muscle and joint pain. Dr. Feely served as the AAO's 2010-11 president. He currently serves as the president of the Foundation for Osteopathic Research and Continuous Education, and he serves on the boards of the American Osteopathic Foundation and The Osteopathic Cranial Academy Foundation.

Registration Fees	Before Oct. 15	On or after Oct. 16
Academy member in practice*	\$700	\$900
Resident or intern member	\$500	\$700
Student member	\$300	\$500
Nonmember practicing DO or other health care professional	\$900	\$1,100
Nonmember resident or intern	\$700	\$900
Nonmember student	\$500	\$700

* The AAO's associate members, international affiliates and supporter members are entitled to register at the same fees as full members.

l am a	nracticing	health	care	professional
rama	practicing	nearth	Care	professional.

□ I am a resident or intern.

□ I am an osteopathic or allopathic medical student.

The AAO accepts check, Visa, MasterCard and Discover payments in U.S. dollars. The AAO does not accept American Express.

Credit card No.: ______Cardholder's name: _______ Expiration date: _______ 3-digit CVV No.: ______

......

Billing address (if different): _____

I hereby authorize the American Academy of Osteopathy to charge the above credit card for the amount of the course registration.

Signature: ____

Click here to view the **AAO's cancellation and refund policy.**

Click here to view the AAO's photo release statement.



Register online at www.academyofosteopathy.org, or submit this registration form and your payment by email to GWatts@academyofosteopathy.org; by mail to the American Academy of Osteopathy, 3500 DePauw Blvd., Suite 1100, Indianapolis, IN 46268-1136; or by fax at (317) 879-0563.



The Incidence of Somatic Dysfunction in Patients With Sudden Onset Atraumatic Neck Pain: A Retrospective Case Note Study

Karen Teten Snider, DO, FAAO; Amrien Ghouse, OMS VI; Sheri Shiyi He, BS; and Vanessa K. Pazdernik, MS

Abstract

Background: Sudden onset atraumatic neck pain is a common complaint, but few studies have investigated this condition.

Objective: To determine the incidence of somatic dysfunction in patients with sudden onset atraumatic neck pain and to assess the immediate responses of the patients to osteopathic manipulative treatment (OMT).

Design: Electronic billing records dated between March 2001 and December 2011 from a neuromusculoskeletal medicine and osteopathic manipulative medicine (NMM/OMM) specialty clinic were retrospectively searched for outpatient clinical encounters that were diagnosed with neck pain or neck strain.

Methods: More than 1000 medical records of patients who were diagnosed with neck pain or neck strain were reviewed for sudden onset pain without precipitating trauma. From the 50 records meeting these criteria, data were collected for demographic characteristics, medical history, signs and symptoms, somatic dysfunctions, types of OMT techniques used, and immediate response to OMT.

Results: Of 1092 neck pain clinical encounters, 50 patients reported sudden onset atraumatic neck pain. Of those 50, patient ages ranged from 10 to 85 years (mean 44 years) and 37 (74%) were female. Twenty-nine patients (58%) had pain onset upon waking. Somatic dysfunctions were documented for 43 patients (86%) in the cervical region, 46 (92%) in the thoracic region, 36 (72%) in the rib region, 13 (26%) in the lumbar region, 8 (16%) in the sacral region, 6(12%) in the pelvic region, and 5(10%)in the cranial region. The most commonly documented muscle spasms were the right levator scapulae and left trapezius (9 patients [18%] for both). Of the 50 patients, 47 (94%) received OMT. The most commonly used OMT techniques included myofascial release used on 30 patients (60%); muscle energy on 26 (52%); highvelocity, low-amplitude on 24 (48%); articular technique on 23 (46%); and counterstrain on 14 (28%). Forty-four patients (88%) reported improved symptoms immediately after OMT.

From the A.T. Still University–Kirksville College of Osteopathic Medicine, Missouri (Snider and Ghouse), From A.T. Still University (Pazdernik), and from Chesterfield, Missouri (He)

Financial disclosure: none reported.

Correspondence address: Karen Teten Snider, DO, FAAO Department of Family Medicine, Preventive Medicine and Community Health A.T. Still University–Kirksville College of Osteopathic Medicine 800 W Jefferson St Kirksville, MO 63501 (660) 626-2304 ksnider@atsu.edu

Submitted for publication Dec. 3, 2015; final revision received May 5, 2016; manuscript accepted May 6, 2016.

Conclusions: The incidence of cervical, thoracic, and rib somatic dysfunction in patients presenting with sudden onset atraumatic neck pain was very high. Females were more likely to present to the clinic with these symptoms, and the onset most commonly occurred on waking. Most patients receiving OMT reported immediate improvement of symptoms.

Introduction

Neck pain is a common musculoskeletal problem that has a substantial impact on health care costs and quality of life.¹ Up to 71% of adults experience neck pain in their lives, and the annual prevalence is estimated between 30% and 50%.² Neck pain is a chronic condition in 10% of those affected.³ The socioeconomic cost of neck pain can be significant and may include absenteeism from work, health care costs, disability, compensation, and impact on the daily activities and work of patients and their families.^{1,4}

(continued on page 10)

(continued from page 9)

Most neck pain is nonspecific with only 10% of neck pain having an identifiable cause, such as an infectious, inflammatory, neurologic, or neoplastic cause.⁴ One type of nonspecific neck pain is commonly referred to as a "crick in the neck" and seems to occur without a precipitating event. The pain occurs suddenly after a simple turn of the head or after trivial atraumatic activities. Common historical findings include "I woke up with neck pain" or "It started after I worked at the computer all day." Typical symptoms include unilateral or bilateral pain accompanied by pain with active or passive cervical rotation. Fryer et al⁵ utilized magnetic resonance imaging of the cervical spine to evaluate 5 patients who reported acute sudden onset atraumatic neck pain. For the level of resolution provided by the magnetic resonance imaging, no common source of pain could be identified.⁵

To determine the incidence of somatic dysfunction in patients with sudden onset atraumatic neck pain, the current study retrospectively reviewed the clinical records of patients presenting to a single neuromusculoskeletal medicine and osteopathic manipulative medicine specialty clinic with sudden onset neck pain of atraumatic origin. Patients' medical histories were assessed along with the incidence of somatic dysfunction findings. The documented response to osteopathic manipulative treatment (OMT) also was reviewed to determine if OMT resulted in reduced symptoms.

Methods

Electronic billing records dated between March 2001 and December 2011 from the Gutensohn Neuromusculoskeletal Medicine/ Osteopathic Manipulative Medicine (NMM/OMM) specialty clinic in Kirksville, Missouri, were retrospectively searched for outpatient encounters involving patients aged 10 years and older who were diagnosed with neck pain or neck strain.

The medical records from the identified encounters were reviewed for documentation indicating an initial evaluation for sudden onset of neck pain without a history of associated trauma. For the current study, sudden onset atraumatic pain was defined as *the abrupt transition from a nonpainful to painful state without a traumatic precipitating event*. Clinical encounters were excluded from the current study if the patient had a documented history of prior cervical or thoracic spinal surgery or a history of trauma to the head, neck, thoracic, or shoulder regions within 6 weeks of the onset of the neck pain. Clinical encounters also were excluded from review if the medical records were stored outside of the clinic.

The study was reviewed and approved by the A.T. Still University-Kirksville Institutional Review Board. The medical records reviewed in the current study included medical histories that were handwritten by patients and medical histories and clinical encounters that were handwritten by the treating physicians or dictated then transcribed. Medical records reviewed for the current study included patients from 22 different attending and resident physicians.

The following data were collected from the medical records:

- total number of OMM encounters related to the neck pain episode
- patient demographics (sex, age, race, ethnicity)
- medical history
- neck pain signs and symptoms
- somatic dysfunctions
- types of OMT used
- immediate response to OMT
- medications prescribed

Information about medical history included history of neck trauma more than 6 weeks prior to onset of neck pain, chronic neck pain, cervical degenerative joint disease, cervical degenerative disc disease, previous episodes of similar neck pain, previous evaluation for current neck pain episode, and onset of neck pain prior to the initial OMM visit.

(continued on page 13)

CONTINUING MEDICAL EDUCATION QUIZ

The purpose of the September 2016 quiz—found on page 20—is to provide a convenient means of self-assessing your comprehension of the scientific content in the article "The Incidence of Somatic Dysfunction in Patients With Sudden Onset Atraumatic Neck Pain: A Retrospective Case Note Study" by Karen Teten Snider, DO, FAAO; Amrien Ghouse, OMS VI; Sheri Shiyi He, BS; and Vanessa K. Pazdernik, MS.

Be sure to answer each question in the quiz. The correct answers will be published in the next issue of the *AAOJ*.

To apply for 2 credits of AOA Category 2-B continuing medical education, fill out the form on page 20 and submit it to the American Academy of Osteopathy. The AAO will note that you submitted the form and forward your results to the American Osteopathic Association's Division of Continuing Medical Education for documentation.

You must score a 75% or higher on the quiz to receive CME credit.

CRANIAL APPROACH OF BERYL E. ARBUCKLE, DO

Dec. 2-4, 2016 • Midwestern University/Arizona College of Osteopathic Medicine in Glendale

This course traces the cranial approaches of William Garner Sutherland, DO; Beryl E. Arbuckle, DO; and Robert Fulford, DO. Dr. Arbuckle was one of Dr. Sutherland's earliest students, studying with him before he taught using primary respiration as a therapeutic force. Diagnosis was performed by palpating the position of the cranial bones and motion testing. In her work with children, Dr. Arbuckle mostly used direct techniques.

Dr. Arbuckle was able to attend hundreds of autopsies, mostly on pediatric neurological cases. She observed regularly arranged fibers in the dura that she called "stress fibers." She also noted that the skull was reinforced in certain places, which she called buttresses. The stress bands and buttresses are used in both diagnosis and treatment.

Attendees will explore the embryology of the head; motion test the sphenobasilar suture, the cranial base, the face, the buttresses, the cranial-cervical junction, and the sacrum; unlock the bony skull and the membranes; work with stress bands; and explore the significance of thoracic respiration.

Continuing Medical Education

24 credits of NMM-specific AOA Category 1-A CME anticipated.

Course Times

Friday and Saturday from 8 a.m. to 6 p.m. Sunday from 8 a.m. to 4 p.m.

Meal Information

Breakfast and lunch will be provided each day. Contact AAO Event Planner Gennie Watts with special dietary needs at (317) 879-1881, ext. 220, or GWatts@academyofosteopathy.org.

Course Location

Midwestern University/Arizona College of Osteopathic Medicine Agave Hall, OMT Lab 101

19555 N. 59th Ave., Glendale, AZ 85308

Travel Arrangements

Contact Tina Callahan of Globally Yours Travel at (800) 274-5975 or globallyyourstravel@cox.net.

_		_
Regio	stratic	n Form
incgi:	in acie	

Cranial Approach of Beryl Arbuckle, DO

Dec. 2-4, 2016

Cours A 1994 A.T. St Osteo DO, se reside medic throug Colleg Athen neuro medic
 Dr. Lo

Course Director

A 1994 graduate of what is now the A.T. Still University–Kirksville College of Osteopathic Medicine, **Kenneth J. Lossing, DO,** served an internship and combined residency in neuromusculoskeletal medicine and family practice through the Ohio University Heritage College of Osteopathic Medicine in Athens. He is board certified in both neuromusculoskeletal medicine and family medicine.

Dr. Lossing studied visceral manipulation

with Jean-Pierre Barral, DO (France). An internationally recognized lecturer, Dr. Lossing contributed to the second and third editions of the American Osteopathic Association's *Foundations of Osteopathic Medicine* textbook.

As the AAO's 2014-15 president, Dr. Lossing was featured in a segment of "American Health Front!" that focused on osteopathic manipulative medicine.

Dr. Lossing and his wife, Margret Klein, OA, run a private practice in San Rafael, California.

Registration Fees	On or before Sept. 30	Oct. 1 through Nov. 26	On or after Nov. 27
Academy member in practice*	\$910	\$960	\$1,160
Resident or intern member	\$710	\$760	\$960
Student member	\$510	\$560	\$760
Nonmember practicing DO or other health care professional	\$1,110	\$1,160	\$1,360
Nonmember resident or intern	\$910	\$960	\$1,160
Nonmember student	\$710	\$760	\$960

* The AAO's associate members, international affiliates and supporter members are entitled to register at the same fees as full members.

- □ I am a practicing health care professional.
- □ I am a resident or intern.
- □ I am an osteopathic or allopathic medical student.

The AAO accepts check, Visa, MasterCard and Discover payments in U.S. dollars. The AAO does not accept American Express.

Name:	/	40A No.:	Credit card No.:	
Nickname for badge:			Cardholder's name:	
Street address:			Expiration date:	3-digit CVV No.:
			Billing address (if different):	
City:	State:	ZIP:		
Phone:	Fax:		I hereby authorize the American Academy o credit card for the amount of the course reg	f Osteopathy to charge the above gistration.
Email:			Signature:	

Click here to view the **AAO's cancellation and refund policy.**

Click here to view the AAO's photo release statement.



Register online at www.academyofosteopathy.org, or submit this registration form and your payment by email to GWatts@academyofosteopathy.org; by mail to the American Academy of Osteopathy, 3500 DePauw Blvd., Suite 1100, Indianapolis, IN 46268-1136; or by fax at (317) 879-0563.



Table 1. Demographic characteristics, medical history, and signs and symptoms of patients with sudden onset atraumatic neck pain (N=50)

Outcome variable	Mean (Range) or No. (%)	Outcome variable
Demographic characteristics		Signs and symptoms
Sex, female	37 (74)	Context of pain onset
Age, y	44 (10 to 85)	On waking
Race		After specific a
White	47 (94)	Unspecified atr
Not specified	2 (4)	Location of pain ^a
Other	1(2)	Right neck
Ethnicity		Left neck
Non-Hispanic	49 (98)	Right shoulder
Not specified	1(2)	Left shoulder
Medical history		Right thoracic
Neck trauma history		Left thoracic pa
Positive trauma > 6 weeks prior to pain onset	2 (4)	Right nonspeci
Negative trauma	39 (78)	Generalized
Unspecified	9 (18)	Unspecified
Chronic neck pain history		Radiation of pain
Positive chronic pain	4 (8)	To head
Negative chronic pain	33 (66)	To shoulders
Unspecified	13 (26)	To arms
Cervical DJD history		To chest
Positive DJD	1(2)	No radiating pa
Negative DJD	32 (64)	Not specified
Unspecified	17 (34)	Character of pain ^a
Cervical DDD history		Stiffness
Positive DDD	1(2)	Tightness
Negative DDD	29 (58)	Ache
Unspecified	20 (40)	Sharp or stabb
Previous episodes of similar pain		Burning
Yes	11 (22)	Not specified
No	14 (28)	Painful active range of
Unspecified	25 (50)	Right rotation
Previous evaluation for current pain episode		Left rotation
Physician outside the OMM clinic	5 (10)	Right sidebend
No other evaluation	32 (64)	Left sidebendir
Not specified	13 (26)	Extension
Onset of pain prior to initial OMM visit		Generalized
≤ 7 days	34 (68)	Unspecified
8-14 days	7 (14)	Progression of pain sir
15-29 days	5 (10)	Improved
1-2 months	2 (4)	Unchanged
> 6 months	1(2)	Worsened
Exact time unspecified but < 6 weeks	1(2)	Unspecified

Outcome variable	Mean (Range) or No. (%)
Signs and symptoms	
Context of pain onset	
On waking	29 (58)
After specific atraumatic activity	13 (26)
Unspecified atraumatic onset	8 (16)
Location of pain ^a	
Right neck	19 (38)
Left neck	18 (36)
Right shoulder	6 (12)
Left shoulder	5 (10)
Right thoracic pain	1(2)
Left thoracic pain	1(2)
Right nonspecific back pain	5 (10)
Left nonspecific back pain	4 (8)
Generalized	16 (32)
Unspecified	1(2)
Radiation of pain	
To head	4 (8)
To shoulders	4 (8)
To arms	2 (4)
To chest	1(2)
No radiating pain	12 (24)
Not specified	27 (45)
Character of pain ^a	
Stiffness	15 (30)
Tightness	5 (10)
Ache	6 (12)
Sharp or stabbing	4 (8)
Burning	1(2)
Not specified	23 (46)
Painful active range of motion ^a	
Right rotation	12 (24)
Left rotation	11 (22)
Right sidebending	o (o)
Left sidebending	1(2)
Extension	1(2)
Generalized	13 (26)
Unspecified	21 (42)
Progression of pain since onset	
Improved	10 (20)
Unchanged	6 (12)
Worsened	9 (18)

Abbreviations: DDD, degenerative disc disease; DJD, degenerative joint disease; OMM, osteopathic manipulative medicine.

^a Patients may have specified 1 or more of the listed signs or symptoms.

25 (50)

(continued from page 10)

For signs and symptoms, information was collected about the context of pain onset, location of pain, radiation of pain away from the site, character of the pain, presence of painful active range of motion, and progression of pain since onset.

Somatic dysfunction assessments along with the number and type of individual somatic dysfunction findings were collected from the physical examination portion of the initial clinical encounter documentation for segmental vertebral dysfunction and tender points in the head, cervical, thoracic, and rib regions along with specific findings of muscular spasms or tender points. Additionally, documentation of somatic dysfunction assessments of the lumbar, pelvic, sacral, and cranial regions was collected from the assessment portion of the clinical encounter. Number and type of somatic dysfunction findings were not collected for lumbar, pelvic, sacral, and cranial regions.

Information regarding types of OMT techniques, if used, immediate response to OMT, and medications prescribed was collected from the plan portion of the initial clinical encounter. Pain intensity after treatment was not documented in the medical records reviewed; therefore pain intensity was not collected in the current study.

Data analysis

The data for patient demographics, medical history, signs and symptoms, somatic dysfunctions, types of OMT techniques used, immediate response to OMT, and medications prescribed were tabulated. The frequency of patients having at least 1 vertebral somatic dysfunction in both the C3-C5 and T2-T5 regions was assessed. Fisher exact test and relative risk were used to assess associations between onset of neck pain and somatic dysfunction findings. $P \leq .05$ were considered statistically significant. The data were analyzed using SAS statistical software (version 9.3, SAS Institute, Inc.).

Results

Of the 1092 outpatient clinical encounters reviewed with neck pain or neck strain, 50 clinical encounters (5%) were documented as having an initial evaluation for sudden onset atraumatic neck pain without history of spinal surgery or recent trauma. The 50 encounters included 50 individual patients. Of those 50 patients,

(continued on page 15)

Table 2. Incidence and frequency of documentation of cervical^a and thoracic^b segmental vertebral somatic dysfunction (N=50)

Vertebral Level	Somatic Dysfunction, No. (%)										
	Total	ESIRI	ESrRr	Extended	FSIRI	FSrRr	Flexed	SIRI	SrRr	NSIRr ^c	NSrRI ^c
C2	17 (34)	3(6)	8 (16)	o (o)	1(2)	2 (4)	0(0)	0(0)	3(6)	-	-
C3	25 (50)	5 (10)	9 (18)	1(2)	3(6)	1(2)	0(0)	2(4)	4 (8)	-	-
C4	21 (42)	3(6)	6 (12)	1(2)	5 (10)	3(6)	0(0)	o (o)	3(6)	-	-
C5	22 (44)	5 (10)	5 (10)	1(2)	4 (8)	4 (8)	0(0)	o (o)	3(6)	-	-
C6	15 (28)	1(2)	3(6)	o (o)	2(4)	4 (8)	o (o)	o (o)	5 (10)	-	-
C7	11 (22)	4 (8)	1(2)	o (o)	3(6)	o(o)	0(0)	o (o)	3(6)	-	-
T1	14 (28)	o (o)	o(o)	o (o)	5 (10)	3(6)	2 (4)	o (o)	2(4)	1(2)	2(4)
T2	29 (58)	1(2)	1(2)	o (o)	3(6)	3(6)	1(2)	o (o)	0(0)	9 (18)	11 (22)
Т3	21 (42)	1(2)	o(o)	o (o)	o(o)	1(2)	o (o)	o (o)	0(0)	10 (20)	9 (18)
Т4	23 (46)	o(o)	o(o)	o (o)	1(2)	1(2)	0(0)	o (o)	0(0)	11 (22)	10 (20)
T5	20 (40)	1(2)	1(2)	1(2)	o(o)	2 (4)	0(0)	o (o)	0(0)	7 (14)	8 (16)
Т6	13 (26)	1(2)	o(o)	o (o)	o(o)	1(2)	0(0)	o (o)	0(0)	6 (12)	5 (10)
T7	9 (18)	o(o)	o(o)	o (o)	o(o)	2 (4)	0(0)	o (o)	0(0)	4 (8)	3(6)
Т8	6 (12)	o (o)	o(o)	o (o)	1(2)	o(o)	o (o)	1(2)	o (o)	o(o)	4(8)
Т9	5 (10) ^d	3(6)	o(o)	o (o)	o(o)	o(o)	0(0)	o (o)	0(0)	1(2)	2 (4)
T10	1(2)	0(0)	0(0)	o (o)	o (o)	o (o)	0(0)	0(0)	0(0)	1(2)	o (o)

Abbreviations: ESIRI, extended, sidebent left, rotated left; ESrRr, extended, sidebent right, rotated right; FSIRI, flexed, sidebent left, rotated left; FSrRr, flexed, sidebent right, rotated right; NSIRr, neutral, sidebent left, rotated right; NSrRI, neutral, sidebent right, rotated left; SIRI, sidebent left, rotated left; SrRr, sidebent right, rotated right.

^a Findings for C1 were limited to 1 patient (2%) with C1 rotated right. ^bNo segmental vertebral somatic dysfunction was documented for T11-T12.

^c Only assessed for thoracic segments.

^d One patient had T9 documented as both NSIRr and NSrRI.

Table :	3. Incidence and freque	ency of documentatio	on of anterior and	posterior cervical	and thoracic ^a tender	points (N=50)
Tubic ,	j. mendernee and ji equi	cincy of abcamentatic	ni oj uncerior una	posterior cervicar	und thoracic tenaci	

Vertebral Level	Somatic Dysfunction No. (%)	Anterior Right Tender Points, No. (%)	Anterior Left Tender Points, No. (%)	Posterior Right Tender Points, No. (%)	Posterior Left Tender Points, No. (%)
C1	3 (6)	2 (4)	o (o)	1(2)	0(0)
C2	1(2)	o (o)	o (o)	1(2)	o (o)
С3	1(2)	o (o)	o (o)	1(2)	o (o)
C4	o (o)	o (o)	o (o)	o (o)	o (o)
C5	1(2)	o (o)	o (o)	1(2)	o (o)
C6	1(2)	1(2)	o (o)	o (o)	o (o)
C7	3 (6)	1(2)	2 (4)	o (o)	o (o)
T1	o (o)	o (o)	o (o)	o (o)	o (o)
T2	o (o)	0(0)	o (o)	o (o)	o (o)
Т3	1(2)	o (o)	o (o)	1(2)	o (o)
T4	1(2)	o (o)	o (o)	1(2)	o (o)

^a No tender points were documented for T5-T12.

Rib Level	Somatic Dysfunction,	Inhaled, ^b	Exhaled, $^{\circ}$	Posterior Tender	Rib Level	Somatic Dysfunction,	Inhaled, $^{\mathrm{b}}$	Exhaled, ^c	Posterior Tender
	No. (%) No. (%) No. (%) Points, ^d No. (%)	Points, ^d No. (%)		No. (%)	No. (%)	No. (%)	Points, ^d No. (%)		
Rib 1					Rib 7				
Right	16 (32)	15 (30)	1(2)	o (o)	Right	1(2)	1(2)	o (o)	o (o)
Left	16 (32)	15 (30)	1(2)	1(2)	Left	o (o)	0 (0)	0 (0)	0(0)
Rib 2					Rib 8				
Right	6 (12)	5 (10)	1(2)	o (o)	Right	o (o)	0(0)	o (o)	o (o)
Left	1(2)	1(2)	o (o)	o (o)	Left	2 (4)	2 (4)	1(2)	o (o)
Rib 3					Rib 9				
Right	5 (10)	4(8)	1(2)	o (o)	Right	o (o)	0(0)	o (o)	o (o)
Left	4 (8)	2 (4)	2(4)	o (o)	Left	1(2)	1(2)	0(0)	o (o)
Rib 4					Rib 10				
Right	6 (12)	3(6)	2 (4)	1(2)	Right	o (o)	o (o)	o (o)	o (o)
Left	4 (8)	2 (4)	2 (4)	o (o)	Left	1(2)	1(2)	o (o)	o (o)
Rib 5					^a No rib s	omatic dysfunct	tions were do	cumented for r	ibs 11-12, and no
Right	3(6)	2 (4)	1(2)	1(2)	anterior	rib tender point	s were docum	ented.	,
Left Rib 6	2 (4)	1(2)	2 (4)	0(0)	^b Includes rib somatic dysfunction documented as elevated without mention of a tender point.				
Right	3(6)	2 (4)	1(2)	1(2)	^c Include	s rib somatic dys	function docu	imented as dep	pressed without
Left	1(2)	1(2)	o (o)	o (o)	mention	of a tender polr	IL.		

^d Includes rib somatic dysfunction documented as a depressed rib tender point.

(continued from page 13)

47 (94%) had 1 clinical encounter related to the neck pain episode, 2 (4%) had 2 clinical encounters for the same episode of neck pain, and 1 (2%) had 3 clinical encounters for the same episode of neck pain. See *Table 1* for detailed results of demographic characteristics, medical history, and signs and symptoms of patients presenting with sudden onset atraumatic neck pain.

Somatic dysfunction was documented at the occipitoatlantal (OA) joint for 27 patients (54%). The OA joint flexed, sidebent right, and rotated left was most frequently documented (11 [22%] patients). Other documented OA diagnoses included extended, sidebent right, and rotated left (5 [10%] patients); flexed, sidebent left, and rotated right (3 [6%] patients); extended, sidebent left, and rotated right (3 [6%] patients); neutral, sidebent right, and rotated left (2 [4%] patients); neutral, sidebent left, and rotated right (2 [4%] patients); and extended (1 [2%] patient). Only 1 patient (2%) had documented dysfunction at C1 (rotated right).

The incidence and frequency of documentation of C2-C7 vertebral somatic dysfunction findings and T1-T10 vertebral somatic dysfunction findings are presented in *Table 2*. Forty-three patients (86%) had at least 1 cervical vertebral or tender point finding documented, with a range findings of 0 to 9 and a mean (SD) of 3.0 (1.9). Forty-six patients (92%) had at least 1 thoracic vertebral or tender point finding documented, with a range of 0 to 10 findings and a mean (SD) of 2.9 (2.2). The most common vertebral somatic dysfunctions were at C3 (25 [50%] patients) and at T2 (29 [58%] patients). Thirty patients (60%) had at least 1 vertebral somatic dysfunction in both the C3-C5 and T2-T5 regions.

The incidence and frequency of documentation of cervical and thoracic tender points is presented in *Table 3*. Five patients (10%) had at least 1 cervical tender point documented, and 1 patient (2%) had at least 1 thoracic tender point documented. The most common cervical tender points were anterior right C1 (2 [4%] patients) and anterior left C7 (2 [4%] patients). The only thoracic tender points documented, right posterior T3 and T4, were in the same patient (1 [2%]). The PC1 inion tender point was documented in 1 patient (2%).

Table 4 lists the incidence and frequency of documentation of rib somatic dysfunction, including tender points. Thirty-six patients (72%) had at least 1 rib dysfunction or tender point documented, with a range of 0 to 7 findings and a mean (SD) of 1.6 (1.7). Inhalation dysfunction was more frequent than exhalation dysfunction, and inhalation of rib 1 was the most frequent finding (15 [30%] on the right and 15 [30%] on the left). Two patients (4%) had 1 or more rib tender points documented.

Table 5. Incidence and frequency of documentation of muscular spasms and tender points (N=50)

Location of Spasm or Tender Poin	t Incidence, No. (%)
Posterior	
Right trapeziu	ıs 7 (14)
Left trapeziu	ıs 9 (18)
Right levator scapula	e 9 (18)
Left levator scapula	e 7 (14)
Right rhomboid	ls 2 (4)
Left rhomboid	ls 2 (4)
Right post scalene	es 5 (10)
Left post scalene	es 5 (10)
Right omohyoi	d 1(2)
Left omohyoi	d o (o)
Right paraspina	al 8 (16)
Left paraspina	al 8 (16)
Right suboccipita	al 1(2)
Left suboccipita	al o (o)
Anterior	
Right anterior scalen	e 2 (4)
Left anterior scalen	e 1(2)
Right middle scalen	e 1(2)
Left middle scalen	e 2 (4)
Right pectoralis minc	or 1(2)
Left pectoralis mind	or o (o)

The incidence and frequency of documented muscular spasms or tender points is presented in *Table 5*. Twenty-eight patients (56%) had at least 1 muscular spasm or tender point documented. The most commonly documented muscle spasms were the trapezius (7 [14%] on the right and 9 [18%] on the left), the levator scapulae (9 [18%] on the right and 7 [14%] on the left) and paraspinal muscles (8 [16%] on the right and 8 [16%] on the left).

Lumbar, sacral, and pelvic somatic dysfunction assessments were documented for 13 (26%), 8 (16%), and 6 (12%) patients, respectively. Cranial somatic dysfunction, excluding the OA joint, was documented for 5 patients (10%).

Twenty-nine patients (58%) had onset of pain on waking, which was negatively associated with right-sided posterior cervical tender points (P=.02). Those patients whose onset of pain did not occur on waking were 15 times more likely to have right posterior cervical tender points than those who reported onset of pain on waking. Onset of pain was not related to any other documented somatic dysfunctions, including tender points or muscle spasms.

Sutherland Cranial Teaching Foundation Upcoming Courses

SCTF Continuing Studies Course: The Eye

October 7-9, 2016



UNE-COM Alfond Center for Health Sciences Biddeford, ME Course Director: Michael Burruano, D.O., F.A.C.

Course cost: \$700 if paid by Aug. 31 \$775 from Sept. 1 on.

October 7, 2016: Registration 11:00 – 11:55am Course begins promptly at 12:00pm October 8, 2016: 8:00am – 5:30pm

October 9, 2016: 9:00am – 12:30pm

Visit our website for enrollment forms and course details: www.sctf.com Contact: Joy Cunningham 907-868-3372 Email: jcunningham4715@yahoo.com

(continued from page 15)

Of the 50 patients, 47 (94%) received OMT for the neck pain. Of the 47 patients who received OMT, myofascial release was the most commonly documented OMT technique. It was used on 30 patients (64%). Muscle energy was used on 26 patients (55%); high-velocity, low-amplitude on 24 (51%); articular technique on 23 (49%); counterstrain on 14 (30%); facilitated positional release on 10 (21%); balanced ligamentous tension or indirect on 9 (19%); percussion hammer on 5 (11%); Still technique on 5 (11%); osteopathic cranial manipulative medicine on 4 (9%); ligamentous articular strain on 4 (9%); neurofascial release on 3 (6%); and soft tissue on 3 (6%). The OMT techniques used on 5 patients (11%) were unspecified.

Of the 47 patients who received OMT, the response was documented as immediate pain improvement for 44 patients (94%) and was unchanged for 1 (2%); posttreatment response was not specified for 2 patients (4%). Medications prescribed included muscle relaxants for 4 patients (8%), nonsteroidal anti-inflammatory medications for 4 patients (8%), and other medications for 14 patients (28%). The other medications may or may not have been related to the neck pain complaint. No medications were prescribed for 28 patients (56%), and 1 patient (2%) received a trigger point injection.

Comment

The current study indicated that 74% of patients seeking NMM/ OMM specialty care for sudden onset atraumatic neck pain were women. A clinical review found that women were more likely than men to visit health care practitioners for nondisabling neck pain.² A recent study also found that women sought treatment for sudden onset nonspecific neck pain more frequently than men.⁶ However, because men are less likely to utilize health care services,^{7,8} these findings do not necessarily reflect the incidence of sudden onset atraumatic neck pain between women and men.

The current study found that only 8% of patients presenting with sudden onset atraumatic neck pain reported chronic neck pain with 2% reporting chronic neck conditions, such as degenerative disc disease. However, over 20% had documented cases of similar neck pain in the past. Leaver et al,⁶ who prospectively investigated acute episodes of nonspecific neck pain, found that 63% of patients reported previous episodes of neck pain. In the current study, information about medical history was limited to that documented by the treating physician; therefore, the incidence of previous episodes of similar neck pain neck pain may have been higher.

In a study by Vasseljen et al,⁹ most patients had significant improvement in pain with or without treatment within 1 month of onset of acute neck pain. In addition, the authors found that those with higher pain intensity were more likely to seek treatment.⁹ Pain intensity data was not collected in the current study because the kind of neck pain studied commonly has 2 levels of pain, one that occurs at rest and a higher pain level that occurs with cervical rotation. Future prospective studies investigating sudden onset atraumatic neck pain should include assessment of pain levels with various cervical ranges of motion.

In the current study, 60% of patients had vertebral somatic dysfunction within C3-C5 and T2-T5. Numerous studies have identified a benefit from treating thoracic dysfunction in the presence of nonspecific neck pain.¹⁰ In one study,¹¹ neck pain patients were randomly assigned to electro/thermal therapy or to electro/thermal therapy plus a seated thoracic thrust manipulation each week for 3 weeks. Those patients receiving the thoracic manipulation had statistically significant improvement in pain (P<.001) and cervical range of motion (P < .05).¹¹ The current study found that somatic dysfunction was common in both the cervical and thoracic regions, suggesting that OMT should be focused in both regions. In a study by Masaracchio et al,¹² mechanical neck pain patients were randomly assigned to combined treatment of the cervical and thoracic spine or to treatment of the cervical spine alone. Their results indicated that the treatment of both regions had better short-term relief of pain than treatment of the cervical spine alone (P < .001).¹²

(continued on page 18)

FULFORD'S ADVANCED PERCUSSION HAMMER

Dec. 2-4, 2016 • University of North Texas Health Science Center **Texas College of Osteopathic Medicine in Fort Worth**

Course Director

When Richard W. Koss. DO. completed

his undergraduate degree at Springfield

College in Massachusetts, he planned to teach physical education, but an

In 1982, Dr. Koss graduated from what is now the A.T. Still University-Kirksville

his focus to osteopathic medicine.

Based on the life exploration, philosophy and osteopathic practice of Robert C. Fulford, DO, this course builds on principles addressed in the basic course. More sensitive and potent concepts and techniques will be presented with supervised practice to assure a proper level of mastery during the course.

Dr. Fulford continued developing new techniques using the percussion hammer, which he pioneered, necessitating the development of an advanced course in the mid 1990s to allow participants to grasp the far-reaching osteopathic healing concepts that were the essence of the old doctor.

This is the same advanced course as taught by Dr. Fulford. An extra day has been added to allow greater study and practice into the thinking and practice of the modern-day osteopathic physician.

Prerequisites

Attendees must have completed a 40-hour introductory cranial course approved by The Osteopathic Cranial Academy and the basic percussion course provided by Dr. Koss.

Continuing Medical Education

22 credits of NMM-specific AOA Category 1-A CME anticipated.

Course Times

Friday and Saturday from 8 a.m. to 6 p.m. Sunday from 9 a.m. to 3 p.m.

Meal Information

Breakfast will be provided Friday and Saturday. Lunch will be provided Friday through Sunday. Contact AAO Event Planner Gennie Watts with special dietary needs at (317) 879-1881, ext. 220, or GWatts@ academyofosteopathy.org.

Course Location

University of North Texas Health Science Center **Texas College of Osteopathic Medicine** 3500 Camp Bowie Blvd., Fort Worth, TX 76107

Travel Arrangements

Contact Tina Callahan of Globally Yours Travel at (800) 274-5975 or globallyyourstravel@cox.net.

Registration Form

Fulford's Advanced Percussion Hammer Dec. 2-4, 2016

I have completed the prerequisites.

Credit card No.: ____ _____ AOA No.: _____ Name: _____ Nickname for badge: ____ Cardholder's name: Street address: _____ Expiration date: ______ 3-digit CVV No.: ______ Billing address (if different): City: _____ State: ____ ZIP: _____ I hereby authorize the American Academy of Osteopathy to charge the above credit card for the amount of the course registration. Phone: _____ Fax: _____ Email: Signature: ____

Click here to view the AAO's cancellation and refund policy.

Click here to view the AAO's photo release statement.



Register online at www.academyofosteopathy.org, or submit this registration form and your payment by email to GWatts@academyofosteopathy.org; by mail to the American Academy of Osteopathy, 3500 DePauw Blvd., Suite 1100, Indianapolis, IN 46268-1136; or by fax at (317) 879-0563.





College of Osteopathic Medicine (ATSU-KCOM) in Missouri, after which he served in the U.S. Air Force Medical Corps for four years as a general medical officer, first at McChord Air Force Base near Tacoma, Washington, and then at Robins Air Force Base in Warner Robins, Georgia.

Dr. Koss first attended a percussion course taught by Robert C. Fulford, DO, in 1987 when Dr. Koss was a resident in osteopathic manipulative medicine at ATSU-KCOM. Two years later, Dr. Fulford invited Dr. Koss to be a table trainer for a percussion course. Dr. Koss continued to assist Dr. Fulford until the latter's death in 1997.

Registration Fees	On or before Sept. 30	Oct. 1 through Nov. 26	On or after Nov. 27
Academy member in practice*	\$814	\$864	\$1,064
Resident or intern member	\$614	\$664	\$864
Student member	\$414	\$464	\$664
Nonmember practicing DO or other health care professional	\$1,014	\$1,064	\$1,264
Nonmember resident or intern	\$814	\$864	\$1,064
Nonmember student	\$614	\$664	\$864

* The AAO's associate members, international affiliates and supporter members are entitled to register at the same fees as full members.

□ I am a practicing health care professional.

□ I am a resident or intern.

I am an osteopathic or allopathic medical student.

The AAO accepts check, Visa, MasterCard and Discover payments in U.S. dollars. The AAO does not accept American Express.

(continued from page 16)

Vertebral tender points were rarely documented for patients in the current study. Because of our retrospective study design, our results may not represent the actual incidence of vertebral tender points because the patients may not have been assessed for vertebral tender points. Documentation of muscular tender points was combined with documentation of muscular spasms during data collection, so we could not determine the incidence of specific muscular tender points. However, the existing documentation of muscular dysfunction suggested that the trapezius, levator scapulae, and paraspinal muscles were most commonly associated with sudden onset atraumatic neck pain. A small study evaluating muscular dysfunction in 15 patients were in the trapezius and levator scapulae.¹³

Although long-term outcomes were not investigated in the current study, 94% of the clinical encounters indicated immediate symptom improvement with OMT. The evidence base of manual therapy intervention for nonspecific neck pain is extensive. A recent systematic review suggested moderate evidence supports using manual therapies to treat patients with acute neck pain, and no evidence suggested that one type of technique was better than another.¹⁴ In the current study, a wide variety of OMT techniques were used to manage neck pain. Future studies could investigate the type of technique or combination of techniques that is most effective for treating patients with atraumatic neck pain.

The current study had several limitations. The retrospective design relied upon physician documentation of all data such as history, physical findings and immediate response to OMT. As such, the data was often nonspecific and may have been incomplete. Therefore, some of the 1092 neck pain encounters reviewed may have met the study's inclusion criteria if documentation of the pain history had been more comprehensive. Another limitation was that the physical examinations performed during the patient encounters did not include a comprehensive assessment for each individual somatic dysfunction reported in the current study. Additionally, because previous studies suggest that the interobserver reliability of specific spinal palpatory findings is fair to poor,^{15,16} correlation of patient symptoms with specific vertebral somatic dysfunctions may have depended on the individual examiner. Future prospective studies should use examiners trained in interobserver reliability to comprehensively assess the correlation of patient history and physical findings with neck pain symptoms.

Conclusion

In the current study, sudden onset atraumatic neck pain was documented in a small percentage of neck pain cases seen in an outpatient NMM/OMM specialty clinic. The actual number of cases may have been higher, however. The incidence of somatic dys-

CLASSIFIED ADVERTISEMENTS

NMM/OMM +1 Positions Available in Corvallis, Oregon

Good Samaritan Regional Medical Center, an active hospital, labor, pediatric and outpatient service, has NMM/OMM +1 residency positions open for 2016-17. Continuity clinics with Oregon State University athletes. Applicants must finish any type of AOAapproved residency by June 2016. Contact Elizabeth Caron, DO, at ecaron@samhealth.org, or call the Graduate Medical Education Department at (541) 768-4906.

3 NMM+1, 3 NMM/FP POSITIONS AVAILABLE IN SUNNY SOUTH FLORIDA

Larkin Hospital in Miami has 3 NMM+1 residencies that begin in July 2016, October 2016 and January 2017. In addition, there are 3 NMM/FP residency opportunities starting in July 2016. Contact Joel D. Stein, DO, FAAO, at (954) 563-2707 or ommdoc@aol.com for more information.

NMM PLUS 1 RESIDENCY PROGRAM IN SOUTHAMPTON NEW YORK

NMM Plus 1 Residency at Southampton Hospital in beautiful Southampton, Long Island. Applications are currently being accepted. If interested, please contact Program Director Lawrence Barnard, DO, Ibarnard@ southamptonhospital.org, Program Coordinator for NMM+1 program Stefania Salzman (631) 726-0409, ext. 126, or Education Department Secretary Karen Roberts at (631) 726-0409.

JOIN BUSY OMT PRACTICE IN CO

Extremely busy established OMT practice in Western CO looking for 1 or 2 physicians to join, then take over the practice. Call Osteopathic Treatment Center at (970) 874-9595.

function was very high in patients presenting with this condition. Consistent with the published literature, women more commonly presented to the clinic for treatment than men, and muscular somatic dysfunction of the trapezius and levator scapulae were common physical findings. Further, OMT resulted in immediate symptomatic improvement in 94% of the patients. The findings from the current study may be useful to guide more comprehensive assessments that can be used in future prospective studies examining the correlation of somatic dysfunction to atraumatic neck pain and the long-term outcomes of OMT.

(continued on page 19)

(continued from page 18)

Acknowledgments

We thank Deborah Goggin, MA, from Research Support at A.T. Still University for her editorial assistance and Emily Burgoon, DO, who contributed to the data collection and research for this project.

References

- Vos T, Flaxman AD, Naghavi M, et al. Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet.* 2012;380(9859):2163-2196. http://www.thelancet.com/ journals/lancet/article/PIIS0140-6736%2812%2961729-2/fulltext. Accessed December 2, 2015.
- Hogg-Johnson S, van der Velde G, Carroll LJ, et al. The burden and determinants of neck pain in the general population: results of the Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders. *J Manipulative Physiol Ther.* 2009;32(2 Suppl):S46-60. http://www.sciencedirect.com/science/article/pii/ S0161475408003370. Accessed December 2, 2015.
- Binder AI. Neck pain [published online first August 4, 2008]. *Clin Evid (Online)*. 2008;2008:1103. http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2907992/. Accessed December 2, 2015.
- Bale A, Newell D. Chiropractic for neck pain: a pilot study examining whether the duration of the pain affects the clinical outcome [published online first September 2, 2005]. *Clin Chiropr.* 2005;8(4):179-188. http://www.sciencedirect.com/science/article/pii/S1479235405000507. Accessed December 2, 2015.
- Fryer G, Adams JH. Magnetic resonance imaging of subjects with acute unilateral neck pain and restricted motion: a prospective case series [published online first January 26, 2011]. *Spine J.* 2011;11(3):171-176. http://www.sciencedirect.com/science/article/ pii/S1529943010014221. Accessed December 2, 2015.
- Leaver AM, Maher CG, McAuley JH, Jull GA, Refshauge KM. Characteristics of a new episode of neck pain [published online first June 18, 2012]. *Man Ther.* 2013;18(3):254-257. http://www.manualthe-rapyjournal.com/article/S1356-689X%2812%2900122-1/abstract. Accessed December 2, 2015.
- Bertakis KD, Azari R, Helms LJ, Callahan EJ, Robbins JA. Gender differences in the utilization of health care services. *J Fam Pract.* 2000;49(2):147-152. http://www.jfponline.com/the-publication/ past-issue-single-view/gender-differences-in-the-utilization-of-healthcare-services/33ae9b4709dda1fdca71aa66c442fec3.html. Accessed December 2, 2015.
- Pinkhasov RM, Wong J, Kashanian J, et al. Are men shortchanged on health? Perspective on health care utilization and health risk behavior in men and women in the United States [published online first February 5, 2010]. *Int J Clin Pract.* 2010;64(4):475-487. http://onlinelibrary.wiley.com/doi/10.1111/j.1742-1241.2009.02290.x/abstract. Accessed December 2, 2015.
- Vasseljen O, Woodhouse A, Bjorngaard JH, Leivseth L. Natural course of acute neck and low back pain in the general population: the HUNT study [published online first April 2, 2013]. *Pain.* 2013;154(8):1237-1244. http://www.sciencedirect.com/science/ article/pii/S0304395913001413. Accessed December 2, 2015.
- 10. Huisman PA, Speksnijder CM, de Wijer A. The effect of thoracic spine manipulation on pain and disability in patients with non-

specific neck pain: a systematic review [published online first January 23, 2013]. *Disabil Rehabil.* 2013;35(20):1677-1685. http://informa-healthcare.com/doi/full/10.3109/09638288.2012.750689. Accessed December 2, 2015.

- Gonzalez-Iglesias J, Fernandez-de-las-Penas C, Cleland JA, del Rosario Gutierrez-Vega M. Thoracic spine manipulation for the management of patients with neck pain: a randomized clinical trial. *J Orthop Sports Phys Ther.* 2009;39(1):20-27. http://www.jospt.org/doi/ pdf/10.2519/jospt.2009.2914. Accessed December 2, 2015.
- Masaracchio M, Cleland JA, Hellman M, Hagins M. Short-term combined effects of thoracic spine thrust manipulation and cervical spine nonthrust manipulation in individuals with mechanical neck pain: a randomized clinical trial. *J Orthop Sports Phys Ther.* 2013;43(3):118-127. http://www.jospt.org/doi/pdf/10.2519/ jospt.2013.4221. Accessed December 2, 2015.
- Munoz-Munoz S, Munoz-Garcia MT, Alburquerque-Sendin F, Arroyo-Morales M, Fernandez-de-las-Penas C. Myofascial trigger points, pain, disability, and sleep quality in individuals with mechanical neck pain. *J Manipulative Physiol Ther.* 2012;35(8):608-613. http://www.sciencedirect.com/science/article/pii/ S0161475412001546. Accessed December 2, 2015.
- Vincent K, Maigne JY, Fischhoff C, Lanlo O, Dagenais S. Systematic review of manual therapies for nonspecific neck pain [published online first November 16, 2012]. *Joint Bone Spine*. 2013;80(5):508-515. http://www.sciencedirect.com/science/article/ pii/S1297319X12002576. Accessed December 2, 2015.
- Nyberg RE, Russell Smith A, Jr. The science of spinal motion palpation: a review and update with implications for assessment and intervention. *J Man Manip Ther.* 2013;21(3):160-167. http://www. ncbi.nlm.nih.gov/pmc/articles/PMC3744849/. Accessed December 2, 2015.
- van Trijffel E, Anderegg Q, Bossuyt PM, Lucas C. Inter-examiner reliability of passive assessment of intervertebral motion in the cervical and lumbar spine: a systematic review [published online first July 1, 2005]. *Man Ther.* 2005;10(4):256-269. http://www.sciencedirect.com/science/article/pii/S1356689X05000640. Accessed December 2, 2015.

Below are the answers to *The AAO Journal's* June 2016 quiz on the article titled "Use of Orthotics to Treat Persistent Low Back Pain After Left Sacrioiliac Joint Fixation: A Case Report" by James A. Lipton, DO, CSP-OMM, FAAO, FAAPMR, FAOCPMR, and 2LT Jochen A. Granja Vasquez MS, MSC, USAR, OMS IV.

- 1. c. According to the article, a successful transacral fixation was described after a failed unilateral fixation.
- **2. e.** Sacroiliac joint fixation surgery can be associated with all of the following: iatrogenic injury, sensorimotor sequellae, hemorrhage, and osseous injury.
- **3. d.** The Young-Burgess Classification pertains to pelvic fractures.
- 4. **e.** The orthotic correction described in this case study involved the use of heel lift and then replacing with custom molded orthotics with built-in assymetrical heel lift and medial wedging.

CONTINUING MEDICAL EDUCATION

This CME Certification of Home Study is intended to document your review of the CME article in this issue of *The AAO Journal* under the criteria for AOA Category 2-B continuing medical education credit.

CME Certification of Home Study

This is to certify that I, _____

(type or print name)

read the following article for AOA CME credit.

Name of article: "The Incidence of Somatic Dysfunction in Patients With Sudden Onset Atraumatic Neck Pain: A Retrospective Case Note Study"

Authors: Karen Teten Snider, DO, FAAO; Amrien Ghouse, OMS VI; Sheri Shiyi He, BS; and Vanessa K. Pazdernik, MS

Publication: *The AAO Journal*, Vol. 26, No. 3, September 2016, pages 9-19

AOA Category 2-B credit may be granted for this article.

00____

(AOA number)

Full name: _____

(type or print name)

Street address: _____

City: ____

State and ZIP code: _____

Signature: _____

Complete the quiz to the right by circling the correct answers. Send your completed answer sheet to the American Academy of Osteopathy. The AAO will forward your results to the American Osteopathic Association. You must answer 75% of the quiz questions correctly to receive CME credit.

Send this page to: American Academy of Osteopathy 3500 DePauw Blvd, Suite 1100 Indianapolis, IN 46268-1136 LGood@academyofosteopathy.org Fax (317) 879-0563

- 1. Which of the data below were collected from medical records?
 - a. Patient demographics
 - b. Medications prescribed
 - c. Type of OMT used
 - d. All of the above
- 2. Which of the following was the most commonly documented technique used?
 - a. Myofascial release
 - b. High-velocity, low-amplitude
 - c. Articular technique
 - d. Counterstrain
- 3. How many patients had vertebral somatic dysfunctions within C3-C5 and T2-T5?
 - a. 40% of patients
 - b. 50% of patients
 - c. 60% of patients
 - d. 70% of patients
- 4. Which of the following was not listed as a limitation of the study?
 - a. The retrospective design relied upon physician documentation of all data such as history, physical findings and immediate response to OMT.
 - b. The physical examinations performed during the patient encounters did not include a comprehensive assessment for each individual somatic dysfunction reported in the current study.
 - c. Because previous studies suggest that the interobserver reliability of specific spinal palpatory findings is fair to poor, correlation of patient symptoms with specific vertebral somatic dysfunctions may have depended on the individual examiner.
 - d. Of the 1092 patients seeking NMM/OMM specialty care for sudden onset atraumatic neck pain, 74% were men.

Answers

See Page 19 for the answers to the *AAOJ's* June 2016 continuing medical education quiz.

Answers to the AAOJ's September 2016 CME quiz will appear in the next issue.

Does Using a Shim Improve the Diagnostic Reliability of the Standing Flexion Test? A Pilot Study

Alexa Paulina Marquez, OMS III, MS; Michelle Yim, OMS IV, MPH; Biana Zelimkhanian, OMS IV, MS; John Quiamas, OMS IV, MS; Rachel E. Davidge, DO; Matthew Siri, DO; and Raymond J. Hruby, DO, MS, FAAODist

Abstract

Context: The standing flexion test (StFT) is a key component in the osteopathic screening exam. Leveling the iliac crests prior to performing the test is thought to improve the reliability of the StFT.

Objective: This pilot study was designed to investigate whether using a shim to level the iliac crests changes the diagnostic reliability of the StFT and to appraise the interexaminer reliability of the StFT between 2 qualified examiners.

Methods: Examiners consisted of 2 osteopathic manipulative medicine predoctoral teaching fellows and 2 first-year osteopathic medical students, all of whom underwent consensus training. Each participant was evaluated by 2 examiners using the StFT, and results were recorded. If iliac crest levels were asymmetrical, a shim was used and a second standing flexion test was performed.

Results: Of the 64 participants in the study, the use of a shim changed the results of the standing flexion test 65.2%, 64.3%, 41.2%, and 64.3% of the time for participants assessed by examiners 1, 2, 3, and 4, respectively. Additional results yielded statistically insignificant interexaminer reliability with moderate reliability (k=0.43) between examiners with higher levels of palpatory experience.

Conclusion: These results suggest that the use of shims to level uneven iliac crests influences the results of the StFT. There is poor interexaminer reliability performing the StFT despite having examiners complete a consensus training session where they reviewed the standardized protocol. However, level of palpatory experience may influence interexaminer reliability.

Introduction

The standing flexion test (StFT) is an integral part of the osteopathic structural exam to assess for pelvic somatic dysfunction. Historically, osteopathic physicians (DOs) are taught that prior to performing the StFT, the iliac crests should be level "to enhance reliability of the standing flexion test."¹ In patients with uneven iliac crest levels, a shim is placed under the patient's heel on the From the Neuromuscular Medicine and Osteopathic Manipulative Medicine Department at the Western University of Health Sciences College of Osteopathic Medicine of the Pacific (WesternU/COMP) in Pomona, California. At the time of the study, Dr Davidge and Dr Siri were osteopathic medical students, and they were predoctoral teaching fellows at WesternU/COMP.

Financial and other disclosures: No financial disclosures reported. Dr Hruby, this article's corresponding author, is the scientific editor emeritus of *The AAO Journal*, and he serves on the journal's editorial review board.

Correspondence address: Raymond J. Hruby, DO, MS, FAAODist Professor and Senior Consultant Department of NMM/OMM Western University of Health Sciences College of Osteopathic Medicine of the Pacific 309 E 2nd St Pomona, CA 91766 (909) 469-5279 rhruby@westernu.edu

Submitted for publication March 31, 2016; final revision received August 17, 2016; manuscript accepted August 17, 2016.

side of the more inferior iliac crest. However, many DOs perform this maneuver with the patient's feet flat on the floor without using shims to adjust for unleveled iliac crests.

Kappler performed a study in which 1 examiner assessed 30 participants and concluded that using shims to adjust for iliac crest levels does not affect the results of this screening maneuver.² With the exception of Kappler's study, which lacked examiner blinding and interexaminer reliability testing, little scientific investigation into the nature of this phenomenon has been done. Therefore, the utility of a shim in the StFT is put to question. This study investigated

(continued on page 22)

(continued from page 21)

whether using a shim to level the iliac crests changes the diagnostic reliability of StFT.

While the StFT has traditionally been taught to all osteopathic medical students at the beginning of their education, it has been widely acknowledged that inconsistent diagnostic interpretation between examiners is also a common problem.

Thus far, the literature demonstrates less than 50% interexaminer reliability of the StFT between osteopathic physicians.³ One factor that may affect the accuracy between examiners is the problem of having poor reliability when using palpatory landmarks during the osteopathic screening exam.⁴⁻⁶ To account for this, the examiners in this study completed consensus training before participants were evaluated. Previous studies have shown that consensus training improves interexaminer reliability⁵ of diagnostic tests.

This pilot study had 2 objectives: to assess whether using a shim to level the iliac crests changes the diagnostic reliability of the StFT and to assess StFT interexaminer reliability among osteopathic medical students. This knowledge will contribute to developing osteopathic diagnostics, and it will lead to improving clinical practice and osteopathic medical student training.

Methods

Participant Selection

A call for volunteer participants was broadcast through emails and announcements in classrooms (eg, osteopathic manipulative medicine laboratory), with no incentive for participation. Potential participants were screened for eligibility by the project investigators and were asked to sign an informed consent to participate in the study.

Participants consisted of 64 volunteers selected from a convenience sample of 124 potential participants from the Western University of Health Sciences student body, faculty, and staff. Participants were males and females 18 years of age or older. Participants who could safely and comfortably bend forward from the hips while standing were eligible for inclusion in the study. Participants were excluded for pregnancy; history of pelvic, femoral, or tibial fracture; lower extremity amputation; hip or knee joint replacement; history of dizziness or vertigo; or any disease that pathologically inhibits lumbar or lower extremity physiologic motion (eg, ankylosing spondylitis).

Consensus Training

All 4 examiners underwent consensus training during a 1-hour session mediated by an osteopathic physician (RJH). Examiners 1 and

Figure. The examiners' consensus training applied this standing flexion test procedure as described by Seffinger et al.⁷

Standing Flexion Test Procedure

- The patient stands with his or her feet comfortably apart and weight evenly distributed to the lower extremities.
- 2. The practitioner sits or kneels behind the patient and places his or her thumbs on the inferior slopes of the patient's PSISs.
- 3. The patient is then asked to bend forward from the waist with arms hanging loosely and to attempt to touch the floor.
- 4. The practitioner observes whether the PSISs move symmetrically (ie, normal motion) or whether 1 PSIS is more superior or ventral than the other at the end of the forward bending motion (ie, positive test result).

2 were first-year osteopathic medical students. Examiners 3 and 4 were predoctoral teaching fellows. Another first-year osteopathic medical student worked as a research coordinator.

The consensus training session entailed locating landmarks, palpating posterior superior iliac spines (PSIS), and standardizing the StFT together on multiple practice participants. The training applied the accepted method of performing the StFT as a step in diagnosing iliosacral somatic dysfunction as described by Seffinger and Hruby.⁷ (See Figure.)

The examiners practiced the StFT until there was an agreement that all examiners were performing the maneuver in the same manner as closely as possible.

Protocol

Each participant was assessed by 2 examiners in a private area. Examiners were blinded to each other. Each examiner performed the StFT with the participant standing and with his or her feet flat on the floor. These results were recorded.

The participant's iliac crest levels were then assessed. If iliac crest levels were asymmetrical, a shim was placed under the inferior iliac crest and a second standing flexion test was performed and results were recorded. The same participant was then directed to the second examiner to perform the same assessment.

Statistical Analysis

Each examiner's data were compared to their own StFT data with and without a shim to determine if placing a shim under the

⁽continued on page 24)

OSTEOPATHIC MANAGEMENT OF CHRONIC PAIN: Addressing Chronic Fatigue Syndrome, Fibromyalgia, Multiple Sclerosis and Neuroinflammation

Jan. 20-22, 2017 • Midwestern University/Arizona College of Osteopathic Medicine in Glendale

Course Director

Bruno J. Chikly, MD, DO (France),

is a graduate of the medical school

and psychiatry. Dr. Chikly also has

the French equivalent of a master's

degree in psychology. He received

at St. Antoine Hospital in Paris,

where his internship in general

medicine included training in endocrinology, surgery, neurology

A growing number of people are diagnosed with chronic fatigue syndrome, fibromyalgia and other chronic pain conditions. These patients need effective, noninvasive treatments that don't aggravate their conditions. This course will provide participants the tools to recognize these conditions and their symptoms and to treat those patients.

The principal treatment will be a blend of Dr. Chikly's lymph and brain techniques. The lymph techniques will include superficial and deeper lymph, mapping and rerouting techniques and some specific viscera work using a lympho-fascia release approach to viscera. The brain techniques will include brainstem, pons, mesencephalon, etc. Specific sacrum intra- or inter-osseous techniques also will be presented.

By the end of the course, participants will be able to identify several chronic pain conditions; describe the comorbidity, known causes, and differential diagnoses of these conditions; explain the importance of the lymphatic system in addressing these conditions; identify megalymphatics in the thorax; and perform lymphatic techniques to release the liver, spleen and small intestines.

Continuing Medical Education

24 credits of NMM-specific AOA Category 1-A CME anticipated.

Course Times

Friday through Sunday from 8 a.m. to 6 p.m.

Meal Information

Breakfast and lunch will be provided each day. Contact AAO Event Planner Gennie Watts with special dietary needs at (317) 879-1881, ext. 220, or GWatts@academyofosteopathy.org.

Course Location

Midwestern University/Arizona College of Osteopathic Medicine Agave Hall, OMT Lab 101

19555 N. 59th Ave., Glendale, AZ 85308

Travel Arrangements

Contact Tina Callahan of Globally Yours Travel at (800) 274-5975 or globallyyourstravel@cox.net.

Registration Form

Osteopathic Management of Chronic Pain

Jan. 20-22, 2017

Name:	AOA No.: O	Ire
Nickname for badge:	(aro
Street address:	E	Exp
	В	Billi
City:	State: ZIP:	hor
Phone:	Fax: c	red
Email:	S	igr

_____0

Click here to view the AAO's cancellation and refund policy.

Click here to view the AAO's photo release statement.



Register online at www.academyofosteopathy.org, or submit this registration form and your payment by email to GWatts@academyofosteopathy.org; by mail to the American Academy of Osteopathy, 3500 DePauw Blvd., Suite 1100, Indianapolis, IN 46268-1136; or by fax at (317) 879-0563.



an honorary DO degree from the European School of Osteopathy in Maidstone, Kent, in the United Kingdom and a PhD in osteopathy from the Royal University Libre of Brussels in Belgium.

Dr. Chikly is an international renowned educator, lecturer and writer. He is the author of the book *Silent Waves: The Theory and Practice* of *Lymph Drainage Therapy*, as well as the creator of a DVD titled *Dissection of the Brain and Spinal Cord*, and he is working on a book about osteopathic manipulation and the brain. He lives in Scottsdale, Arizona, with his wife and partner, Alaya.

Registration Fees	On or before Nov. 19, 2016	Nov. 20, 2016 through Jan. 14, 2017	On or after Jan. 15, 2017
Academy member in practice*	\$970	\$1,020	\$1,220
Resident or intern member	\$770	\$820	\$1,020
Student member	\$570	\$620	\$820
Nonmember practicing DO or other health care professional	\$1,170	\$1,220	\$1,420
Nonmember resident or intern	\$970	\$1,020	\$1,220
Nonmember student	\$770	\$820	\$1,020

* The AAO's associate members, international affiliates and supporter members are entitled to register at the same fees as full members.

- □ I am a practicing health care professional.
- □ I am a resident or intern.
- □ I am an osteopathic or allopathic medical student.

The AAO accepts check, Visa, MasterCard and Discover payments in U.S. dollars. The AAO does not accept American Express.

Credit card No.: _____

Cardholder's name: ____

Expiration date: _______ 3-digit CVV No.: ______

Billing address (if different): _____

hereby authorize the American Academy of Osteopathy to charge the above credit card for the amount of the course registration.

Signature: _

Register online at www.acade

(continued from page 22)

patient's foot made a difference in screening for pelvic somatic dysfunction. Percentage agreement was calculated to evaluate the outcome of the use of shims on the standing flexion test. Cohen's kappa coefficient (κ) and their 95% confidence intervals were calculated to evaluate interrater reliability. K factors out agreement between results due to chance. The κ coefficient ranges from -1 to 1 where 0 indicates an agreement no better than chance and positive values signify agreement better than results due to chance alone. The authors followed the guidelines in *Table 1* suggested by Landis and Koch⁸ to interpret κ .

Results

Of the 64 participants of both genders who met the inclusion criteria, 46 participants were assessed by examiners 1 and 2 while 18 participants were assessed by examiners 3 and 4. The use of a shim changed the results of the standing flexion test 65.2%, 64.3%, 41.2%, and 64.3% of the time for participants assessed by examiners 1, 2, 3, and 4 respectively. *(See Table 2)*

In evaluating the data, it was determined that the difference in the number of participants assessed by each set of examiners was not statistically significant.

Table 1. Guidelines by Landis and Koch for interpreting κ.

Value of κ	Strength of agreement	
0.0-0.20	Slight	
0.21-0.40	Fair	
0.41-0.60	Moderate	
0.61-0.80	Substantial	
0.81-1.00	Almost perfect	

Table 2. The effect of shims on the standing flexion test.

Examiner	n	Uneven iliac crest	Different StFT result with shim	Percentage
1	46	23	15	65.7%
2	46	28	18	64.3%
3	18	17	7	41.2%
4	18	14	9	64.3%

Table 3. Interrater reliability between examiners.

Kappa was calculated to determine the level of agreement in using a shim between examiners 1 and 2, (κ_1 =0.16) and examiners 3 and 4 (κ_2 =0.43), which correlates to slight and moderate agreement respectively. Additional results did not indicate significant agreement between the 2 examiners of each group in measuring iliac crest symmetry (κ_1 =0.19, κ_2 =0.37) or in agreement on initial StFT results (κ_1 = 0.29, κ_2 =-0.08).

Discussion

The results of this study indicate that using shims to level uneven iliac crests alters the outcome of the StFT. These results oppose conclusions made by Kappler. This study challenged Kappler's results by having 2 examiners, rather than 1 examiner, assess each study participant. Both examiners found shims to affect the StFT results, suggesting that leveling the iliac crest with a shim may be necessary prior to performing the StFT. One reason for this finding may be that unleveled iliac crests may also suggest that PSIS levels are uneven. Thus, when performing the StFT, structural asymmetry of the iliac crests may alter the degree of excursion of the PSIS when compared to leveled iliac crests.

Additionally, the results of this study indicate poor to moderate interexaminer reliability of the osteopathic structural exam components. This evidence is consistent with previous literature.^{5,9}

Studies also have found that prior training to prepare researchers for a specific study protocol will increase interexaminer reproducibility.^{10,11} This principle was carried into the current study in an attempt to maximize interexaminer reproducibility and to control for the difference in experience between examiners. Thus, consensus training was performed, but it was only performed once prior to initiating data collection.

Despite these efforts to control for interexaminer variability, it is apparent that the consensus training did not have a significant effect on minimizing the variability in the structural assessment and StFT results between examiners.

> Perhaps having multiple sessions of consensus training (ie, prior to each set of data collection) would have helped to reduce inconsistencies and to improve accuracy in following standardized pro-

> > (continued on page 25)

Agreement	Examiners 1 and 2 (κ_1)	Examiners 3 and 4 (κ_2)
Iliac crests level	0.19 (95% Cl -0.13, 0.50)	0.37 (95% CI -0.12, 0.85)
Shim necessity	0.16 (95% Cl -0.14, 0.47)	0.43 (95% Cl 0.05, 0.81)
First standing flexion test	0.29 (95% CI -0.08, 0.67)	-0.08 (95% CI -0.19, 0.03)

(continued from page 24)

tocols and locating anatomical landmarks. It is important that institutions find methods that optimize students' abilities to consistently follow protocol.

This brings into question the amount of variability in protocols used for the same osteopathic diagnostic procedures. For example, some StFT protocols require participants to position their feet shoulder width apart¹² versus comfortably apart.⁷ In support of this, Bogduk¹³ describes the need to establish standardized protocols for diagnostic assessments in osteopathic medicine. If there is poor interexaminer reliability within a standardized protocol, then it suggests that there may be wider variability in diagnoses when using different protocols for a particular screening test.

It is also important to point out that there was a stronger agreement between examiners 3 and 4 when compared to examiners 1 and 2 ($\kappa_2 > \kappa_1$) in using a shim and measuring iliac crest symmetry. This may be attributed to the fact that examiners 3 and 4 have a higher level of palpatory experience when compared to examiners 1 and 2. This suggests that level of training may strengthen consistency in the structural evaluation between examiners. However this data may have been skewed due to the fact that examiners 1 and 2 assessed 28 more participants than examiners 3 and 4.

Study Limitations

One limitation of this study was the small sample size. More studies of this nature, using larger groups of participants, would provide for a higher level of confidence in the results.

Another limitation in this study is the varying levels of experience between examiners. The outcome may have been different if all examiners were of the same training level.

One last factor to take into consideration is that the participants were primarily healthy medical students. It would be interesting to assess whether this study would have had a different outcome on patients with pelvic somatic dysfunctions.

Conclusion

This pilot study demonstrated that the use of shims to level uneven iliac crests influences the results of the StFT.

Poor interexaminer reliability remains an issue despite providing consensus training and reviewing the standardized protocol. However, interexaminer reliability may be strengthened by an examiner's level of palpatory experience and should be explored further.

Additional studies should aim to obtain a greater sample size, to use examiners of the same experience level, to have more standardized

consensus training sessions, and to create more standardized diagnostic protocol. The authors hope this knowledge might contribute to the development of osteopathic diagnostics and lead to improving clinical practice and osteopathic medical student training.

Acknowledgements

The authors thank Dwayne Townsend for data entry and Eric Hurwitz, DC, PhD, for conducting the statistical analysis of the data gathered for this study.

References

- 1. Mitchell FL, Mitchell PK. *The muscle energy manual, Vol. 3: evaluation and treatment of the pelvis and sacrum.* East Lansing, MI: MET Press; 2001:86.
- Kappler RE. Standing flexion test study. AAO Journal. 2000;10(3):29-30.
- Vincent-Smith, B, Gibbons, P. Inter-examiner and intra-examiner reliability of the standing flexion test. *Manual Therapy*. 1999;4(2):87-93. http://www.manualtherapyjournal.com/article/ S1356-689X(99)90173-X/abstract. Accessed January 28, 2016.
- Sutton C, Nono L, Johnston RG, Thomson OP. The effects of experience on the inter-reliability of osteopaths to detect changes in posterior superior iliac spine levels using a hidden heel wedge. *J Bodyw Mov Ther.* 2013;17(2):143-150.
- Degenhardt BF, Snider KT, Snider EJ, Johnson JC. Interobserver reliability of osteopathic palpatory diagnostic tests of the lumbar spine: Improvements from consensus training. *J Am Osteopath Assoc*. 2005;105(10):465-473.
- Stovall BA, Kumar S. Reliability of boney anatomic landmark asymmetry assessment in the lumbopelvic region: Application to osteopathic medical education. *J Am Osteopath Assoc.* 2010;110(11):667-674.
- Seffinger MA, Hruby RJ, Kuchera WA, illustrator. *Evidence-Based* Manual Medicine: A Problem-Oriented Approach. Philadelphia, PA: Saunders Elsevier; 2007: 40-42.
- 8. Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics*. 1977;33:159-174.
- Rajendran D, Gallagher D. The assessment of pelvic landmarks using palpation: A reliability study of undergraduate students. *Int J Osteopath Med.* 2011;14(2):57-60.
- Fryer G, McPherson HC, O'Keefe P. The effect of training on the inter-examiner and intra-examiner reliability of the seated flexion test and assessment of pelvic anatomical landmarks with palpation. *Int J Osteopath Med.* 2009;8(4):131-138.
- Gerwin R, Shannon S, Hong C, Hubbard D, Gervirtz R. Inter-rater reliability in myofascial trigger point examination. *Pain*. 1997;69(1-2):65-73.
- 12. Ehrenfeuchter WC. Screening osteopathic structural examination. In: Chila AG, executive ed. *Foundations of Osteopathic Medicine*. 3rd ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2011:427.
- Bogduk, N. An interview with Nikolai Bogduk. Musculoskeletal medicine, evidence based medicine and the meaning of life. Newsletter of the Australian Association of Musculoskeletal Medicine. 1998:1-4.

American Academy of Osteopathy*

2017 Convocation

March 22-26, 2017 • The Broadmoor • Colorado Springs

Pre-Convocation Courses

March 18-21, 2017 Pre-Convocation course: The Strategic Crossroads of the Body Course director: Kenneth J. Lossing, DO Featured speaker: Jean-Pierre Barral, DO (UK) 32 credits of NMM-specific AOA Category 1-A CME anticipated

March 19-21, 2017 Pre-Convocation course: Brain 1: Palpating and Treating the Brain, Brain Nuclei, White Matter and Spinal Cord Course director: Bruno J. Chikly, MD, DO (France) 24 credits of NMM specific AOA Category 1-A CME anticipated March 19-21, 2017 Pre-Convocation course: Fascial Distortion Model: Axial Spine Course director: Todd A. Capistrant, DO, MHA 24 credits of NMM specific AOA Category 1-A CME anticipated

> March 22, 2017 Pre-Convocation course: Cervical Glymphatics Course director: Frank Willard, PhD 4 credits of NMM specific AOA Category 1-A CME anticipated



2017 Convocation

The Balance Point:

Bringing the Science and Art of Osteopathic Medicine Together

Natalie Ann Nevins, DO, program chair

www.academyofosteopathy.org

AAOJ Submission Checklist

Manuscript Submission

- □ Submission emailed to editoraaoj@gmail.com or mailed on a flash drive or CD to the *AAOJ* managing editor, American Academy of Osteopathy, 3500 DePauw Blvd, Suite 1100, Indianapolis, IN 46268-1136
- Manuscript formatted in Microsoft Word for Windows (.doc, .docx), text document format (.txt), or rich text format (.rtf)

Manuscript Components

- □ Cover letter addressed to the *AAOJ's* scientific editor with any special requests (eg, rapid review) noted and justified
- □ Title page, including the authors' full names, financial and other affiliations, and disclosure of financial support related to the original research or other scholarly endeavor described in the manuscript
- □ "Abstract" (see "Abstract" section in "*AAOJ* Instructions for Contributors" for additional information)
- □ "Methods" section
 - the name of the public registry in which the trial is listed, if applicable
 - ethical standards, therapeutic agents or devices, and statistical methods defined
- □ Four multiple-choice questions for the continuing medical education quiz and brief discussions of the correct answers
- □ Editorial conventions adhered to
 - terms related to osteopathic medicine used in accordance with the *Glossary of Osteopathic Terminology*
 - units of measure given with all laboratory values
 - on first mention, all abbreviations other than measurements placed in parentheses after the full names of the terms, as in "American Academy of Osteopathy (AAO)"
- Numbered references, tables, and figures cited sequentially in the text
 - journal articles and other material cited in the "References" section follow the guidelines described in the most current edition of the AMA Manual of Style: A Guide for Authors and Editors
 - references include direct, open-access URLs to posted, full-text versions of the documents, preferably to digital object identifiers (DOIs) or to the original sources
 - photocopies provided for referenced documents not accessible through URLs
- □ "Acknowledgments" section with a concise, comprehensive list of the contributions made by individuals who do not merit

authorship credit, as well as permission from each individual to be named

□ For manuscripts based on survey data, a copy of the original validated survey and cover letter

Graphic Elements

- □ Graphics formatted as specified in the "Graphic Elements" section of "*AAOJ* Instructions for Contributors"
- □ Graphics as separate graphic files (eg, jpg, tiff, pdf), not included with text
- □ Each graphic element cited in numerical order (eg, *Table 1*, *Table 2* and *Figure 1*, *Figure 2*) with corresponding numerical captions provided in the manuscript
- □ For reprinted or adapted tables, figures, and illustrations, a full bibliographic citation given, providing appropriate attribution

Required Legal Documentation

- □ For reprinted or adapted tables, figures, and illustrations, copyright holders' permission to reprint in the *AAOJ*'s online and print versions, accompanied by photocopies of the original published graphic designs
- □ For photographs in which patients are featured, signed and dated patient model release forms
- □ For named sources of unpublished data and individuals listed in the "Acknowledgments" section, written permission to publish their names in the *AAOJ*
- □ For authors serving in the US military, the armed forces' written approval of the manuscript, as well as military or other institutional disclaimers

Financial Disclosure and Conflict of Interest

Authors are required to disclose all financial and nonfinancial relationships related to the submission's subject matter. All disclosures should be included in the manuscript's title page. See the "Title Page" section of "*AAOJ* Instructions to Contributors" for examples of relationships and affiliations that must be disclosed. Those authors who have no financial or other relationships to disclose must indicate that on the manuscript's title page (eg, "Dr Jones has no conflict of interest or financial disclosure relevant to the topic of the submitted manuscript").

Publication in the JAOA

Please include permission to forward the manuscript to *The Journal* of the American Osteopathic Association if the AAOJ's scientific editor determines that the manuscript would likely benefit osteopathic medicine more if the JAOA agreed to publish it.

Questions? Contact editoraaoj@gmail.com.

Component Societies and Affiliated Organizations Calendar of Upcoming Events

Sept. 16-18, 2016 The Osteopathic Cranial Academy Orofacial Development: Merging OCF and Functional Dentistry Course director: Eric J. Dolgin, DO, FCA Hilton Irvine Hotel in California 22.5 credits of AOA Category 1-A CME anticipated Learn more and register at www.cranialacademy.org.

Sept. 17, 2016 Osteopathy's Promise to Children OMT for Systemic Disorders and Physiological Functions: Cardiopulmonary and Immune Systems Course director: Hollis H. King, DO, PhD, FAAO Osteopathic Center, San Diego & credits of AOA Category 1-A CME anticipated Learn more and register at www.the-promise.org/cme.

Sept. 17, 2016

FIMM (International Federation for Manual/Musculoskeletal Medicine) and the Bulgarian Society of Manual Medicine (BSMM) *New Evidence in Science Related to Manual and Musculoskeletal Medicine* Varna, Bulgaria Learn more and register at fimm-online.com.

Sept. 18-19, 2016

FIMM (International Federation for Manual/Musculoskeletal Medicine) and the Bulgarian Society of Manual Medicine (BSMM) **2nd International Instructors Course** Medical faculty, Varna, Bulgaria Learn more and register at fimm-online.com.

Sept. 18-19, 2016

FIMM (International Federation for Manual/Musculoskeletal Medicine) and the Bulgarian Society of Manual Medicine (BSMM) **Dry Needling and Manual Medicine for Myofascial Pain** Medical faculty, Varna, Bulgaria Learn more and register at fimm-online.com.

Sept. 24-25, 2016 New York Institute of Technology College of Osteopathic Medicine OMT for the Visceral System Course directors: Johannes Mayer, MD, DOM (Germany), and Simeon John Hain, DO Old Westbury, New York 14.25 credits of AOA Category 1-A CME Learn more and register at NYIT.edu.

Sept. 30-Oct. 4, 2016

Michigan State University College of Osteopathic Medicine Craniosacral Techniques: Part II Course director: Barbara J. Briner, DO East Lansing, Michigan 35 credits of AOA Category 1-A CME anticipated Learn more and register at com.msu.edu.

Oct. 7-9, 2016

Rocky Mountain Academy of Osteopathy **Querying the Body Using PRM and Long Fascial Relationships** Course director: R. Paul Lee, DO, FAAO, FCA Rocky Vista University College of Osteopathic Medicine Parker, Colorado 24 credits of AOA Category 1-A CME anticipated Learn more and register at rockymountainaao.wix.com/rockymtnaao.

Oct. 7-9, 2016

Sutherland Cranial Teaching Foundation The Eye Course director: Michael P. Burruano, DO, FCA University of New England College of Osteopathic Medicine Biddeford, Maine Learn more and register at www.sctf.com.

Oct. 8, 2016

Osteopathy's Promise to Children **OMT for Systemic Disorders and Physiological Functions: Gastrointestinal and Nervous Systems** Course director: Hollis H. King, DO, PhD, FAAO Osteopathic Center, San Diego 8 credits of AOA Category 1-A CME anticipated Learn more and register at www.the-promise.org/cme.

Oct. 21-24, 2016

Michigan State University College of Osteopathic Medicine Exercise Prescription as a Complement to Manual Medicine Course director: Mark Bookhout, PT Course faculty: Lisa Chase, PT East Lansing, Michigan 26 credits of AOA Category 1-A CME anticipated Learn more and register at com.msu.edu.

Nov. 4-6, 2016

Osteopathy's Promise to Children **A Sequential Approach to Pediatric Osteopathy** Course director: Mary Anne Morelli Haskell, DO, FACOP Osteopathic Center, San Diego 24 credits of AOA Category 1-A CME anticipated Learn more and register at www.the-promise.org.

Visit www.academyofosteopathy.org for additional listings.